

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

USE OF PHANQUINONE FOR THE TREATMENT OF ALZHEIMER'S DISEASE

TECHNICAL FIELD

The present invention relates to the use of a
5 known compound for the manufacture a pharmaceutical
composition for treatment or prevention of Alzheimer's
disease. Further, the invention relates to a pharma-
ceutical composition for such treatment or prevention.
The invention is also directed to methods of treatment
10 or prevention of Alzheimer's disease.

BACKGROUND OF THE INVENTION

Alzheimer's disease (AD), which is the single
major cause of dementia in adults in industrialized
15 societies, is a degenerative brain disorder character-
ized clinically by a progressive loss of memory, con-
fusion, dementia and ultimately death. Histopathologi-
cally, Alzheimer's disease is characterized by the
presence in the neocortex, especially the hippocampus,
20 of two brain lesions: the neurofibrillary tangles
(NFTs) of paired helical filaments (PHF) in the neurons
and the neuritic (senile) plaques in the extracellular
space. The formation of senile plaques is related to
the appearance of the symptoms and signs of the dis-
25 ease, including amnesia. After the formation of senile
plaque, neurofibrillary tangles are produced in the
neuronal bodies. The formation of neurofibrillary
tangles is related to the worsening of amnesia and of
the other symptoms of dementia.

30 A major component of the senile plaques is amyloid
deposits. Cataract is a disease often occurring
together with Alzheimer's disease. It is located in the
eyes and is also caused by amyloid deposits. Therefore,
the treatment of patients having Alzheimer's disease
35 disclosed in the present description and claims may

equally be used to treat cataract. As used in the present description and claims, the term Alzheimer's disease, therefore, also include the disease cataract.

A main component of the amyloid deposits is a polypeptide referred to herein as A β (Amyloid-beta). A β is normally a soluble component of the cerebrospinal fluid where it is found in concentrations of about 3-5 nm. A β may have 39 to 43 amino acids, typically 40 amino acids, in the mature form and is derived as a proteolytic cleavage product from a cell surface protein called the amyloid precursor protein (APP) (Kang et al. 1987, Nature 325:733-736):

Many studies have shown that A β is toxic in vitro when added directly to neuronal cell cultures (Yankner BA, Duffy LK, Kirschner DA, Science 1990, 250 (4978): 279-282; Koh JY, Yang LL, Cotman CW, Brain Res 1990, 533 (2): 315-320; and Pike CJ, Burdick D, Walencewicz AJ, Glabe CG, Cotman CW, J. Neurosci 1993, 13(4):1676-1687).

The neurotoxicity of A β has been located to be in the peptide sequence between amino acid residues 25 and 35 (A β (25-35)). A β (25-35) induces neuronal cell death equally potent as full length A β (1-40) (Yankner BA, Duffy LK, Kirschner DA, Science 1990, 250 (4978): 279-282). The normal function of A β is not known at present but might be to form cation selective channels across cell membranes (Kawahara M. et al., 1997, Biophysical Journal 73/1, 67-75).

The precipitation of synthetic A β has been shown to be caused by several environmental factors including low pH, high salt concentrations and the presence of metals, e.g. zinc, copper, and mercury (Bush, A.I. et al., 1995, Science 268: 1921-1923). It has been reported that A β itself specifically and saturable binds zinc with a high affinity binding (K_D = 107 nM) at a molar

ratio of 1:1 (zinc: A β) (Bush, A.I. et al., 1994, J. Biol. Chem. 269: 12152-12158). This binding takes place at physiological concentrations of zinc (Bush, A.I. et al., 1994, Science 265: 1464-1467).

5 There is a strong supposition that removal of amyloid deposits from patients suffering from Alzheimer's disease will alleviate the symptoms of Alzheimer's disease. Therefore, several attempts have been made to prepare such a drug, as methods for
10 healing Alzheimer's disease are urgently sought.

International Patent Application, publication No. WO 93/10459, discloses a method for the treatment of Alzheimer's disease by administering a zinc binding agent. As preferred compounds, phytic acid, desferri-
15 oximine, sodium citrate, EDTA, 1,2-diethyl-3-hydroxypyridine-4-one, and 1-hydroxyethyl-3-hydroxy-2-methylpyridine-4-one are mentioned.

German patent application No. DE 39 32 338 discloses the use of an aluminium chelator, such as 8-
20 hydroxy-quinoline, for the treatment of Alzheimer's disease.

U.S. Patent No. 5,373,021 discloses the use of disulfiram and its salts and analogs. According to this patent, the disclosed compounds may be used to reduce
25 neurological damage caused by Alzheimer's disease.

International Patent Application, publication No. WO 98/06403 discloses the use of clioquinol for the manufacture of a pharmaceutical composition for the treatment of Alzheimer's disease.

30 The hitherto known compounds suggested for the treatment of Alzheimer's disease have several drawbacks, which has prevented their widespread use. Most of the compounds are unable to penetrate the blood-brain-barrier and thus cannot readily reach the areas
35 in which the amyloid is deposited. Disulfiram, which

may penetrate the blood-brain-barrier, has the drawback that, when it is combined by a patient with ethyl alcohol, it causes severe adverse reactions, including headaches, nausea, vomiting, sweating, thirst, weakness, and low blood pressure. Cliquinol (5-chloro-7-iodo-8-hydroxyquinoline), which also may penetrate the blood-brain-barrier, has a damning history as it as a side effect causes subacute myelo-optico-neuropathy (SMON).

10 Phanquinone (4,7-phenanthroline-5,6-dione) has hitherto been used for the treatment of various disorders, such as amoebiasis. Phanquinone has been sold by CIBA-GEIGY under the trademark ENTOBEX. In contrast to clioquinol no adverse side effects have been
15 detected when phanquinone is used in the normal dosage range.

In the past an antiamebic pharmaceutical preparation containing both clioquinol and phanquinone has been sold by CIBA GEIGY under the trademark Mexafor.
20 However, the marketing of this preparation was stopped when it was realized that clioquinol caused SMON.

DISCLOSURE OF THE INVENTION

The object of the present invention is to provide
25 a new use of a known compound for the treatment or prevention of Alzheimer's disease. Another object of the present invention is to provide for an increased effect of the treatment of Alzheimer's disease. A further object is to avoid any detrimental side effect
30 of the treatment or prevention of Alzheimer's disease.

According to the present invention the use of phanquinone for the manufacture of a pharmaceutical composition for the treatment or prevention of Alzheimer's disease is provided.

Phanquinone may be administered in any amount efficient for the treatment or pervention of Alzheimer's disease. Preferably, phanquinone may be administered in an amount of 5 mg to 250 mg, and most 5 preferred 10 mg to 50 mg, one to three times daily.

In one embodiment of the invention, a compound, or a mixture of compounds, selected from the group comprising antioxidants, acetylcholine enhancers, trace metals, prosthetic groups and clioquinol, is administered prior to, together with or subsequent to the 10 administration of phanquinone.

The invention also relates to a pharmaceutical composition comprising phanquinone and a compound, or a mixture of compounds, selected from the group comprising 15 antioxidants, acetylcholine enhancers, trace metals, prosthetic groups and clioquinol provided, when clioquinol is selected, that at least one further compound is selected from the group.

The antioxidant is preferably vitamin C, vitamin 20 E, Q10, or combinations thereof. The acetylcholine enhancers are preferable μ 1 agonists or anticholinesterase inhibitors. Preferred anticholinesterase inhibitors are tacrine (trademark: Cognex) and donepezil (trademark: Aricept). The prosthetic group is preferably vitamin B₁₂. 25

In a preferred embodiment of the invention phanquinone and clioquinol are used for the manufacture of the pharmaceutical composition for the treatment or prevention of Alzheimer's disease. Preferably, a 30 combination of phanquinone, clioquinol and vitamin B₁₂ is used for the manufacture of the pharmaceutical composition.

The pharmaceutical composition may be formulated for oral, parenteral or intradermal administration. 35 Further, the pharmaceutical formulation may be formu-

lated as a single pharmaceutical composition or as two or more separate pharmaceutical entities for sequential or substantially simultaneous administration.

The invention also relates to a method of treating a subject having or suspected of having Alzheimer's disease comprising administering to the subject an amount of phanquinone effective to treat or prevent Alzheimer's disease.

10 BRIEF DESCRIPTION OF THE FIGURES

Figure 1 depicts the effect of phanquinone on $A\beta(25-35)$ dose-response in PC12 cells.

Figure 2 depicts the effect of phanquinone in $A\beta(25-35)$ induced toxicity in PC12 cells.

15 Figure 3 depicts the effect of phanquinone and clioquinol on Zn^{2+} -induced $A\beta$ aggregation.

Figure 4 depicts the effect of clioquinol and phanquinone on Cu^{2+} -induced $A\beta$ aggregation.

Figure 5 depicts the NMR spectra of three solutions. Solution a contained vitamin B_{12} (hydroxycobalamin), at a concentration of 2.6 mM. Solution b contained a mixture of 2.6mM vitamin B_{12} and 10 mM clioquinol glucuronide (mole ratio of about 1:4). Solution c contained 10 mM clioquinol glucuronide.

25 Figure 6 depicts the right half of figure 5 expanded for easier comparison of the resonance positions.

Figure 7 depicts the left half of figure 5 expanded for easier comparison of the resonance positions.

DETAILED DESCRIPTION OF THE INVENTION

There is a need for introducing a drug on the market that is efficient in the therapy or the prophylaxis of Alzheimer's disease. According to the present

invention the known drug phanquinone is suggested.

In the examples it is shown that $A\beta(25-35)$ in PC12 cells has a pronounced effect on the redox activity of cells. In the method used in example 2, the redox activity of cells exposed to $1 \mu M A\beta(25-35)$ is reduced to 60% of the normal level (see fig 2, open circles). However, when the $A\beta(25-35)$ -treated cells is treated with $1 \mu g/ml$ phanquinone or more the normal redox activity is restored. This indicate that phanquinone when administered to a mammal such as a human will alleviate the adverse effects of $A\beta$.

While it is not intended to limit the invention to any specific mechanism of action, it is at present believed that it is an antioxidant effect of phanquinone that prevent the effect of $A\beta$ on the cell redox activity.

The antioxidant effect of phanquinone may be improved by co-administering further antioxidant(s) such as vitamin C, vitamin E, vitamin Q10 or combinations thereof.

The presence of senile plaques in the extracellular space may prevent or inhibit the transmission of impulses in the cholinergic nervous system. The functioning of the nervous system is dependent on the mediator acetylcholine. Therefore, compounds enhancing the level of acetylcholine, such as compounds preventing or inhibiting the normal hydrolysis of acetylcholine by acetylcholinesterase, so-called anticholinesterase inhibitors, may be administered together with phanquinone to improve the effect. Preferred anticholinesterase inhibitors are tacrine and donepezil.

In a study reported in the examples, it is shown that phanquinone can reduce the aggregation of $A\beta(1-40)$ induced by Cu^{2+} and Zn^{2+} .

The following proposed mechanism of action of the invention is not intended to limit the invention to said mechanism. At present, applicant believes that phanquinone and $A\beta$ competitively chelate zinc, copper and other heavy metals. Phanquinone has the ability to penetrate the blood-brain-barrier. When phanquinone has captured a heavy metal ion in the extracellular space it moves into the blood and is cleared from the body. As the aggregation or salt of $A\beta$ and the heavy metal ion is in equilibrium with free $A\beta$ and the free heavy metal ion phanquinone may continue to capture free heavy metal ions and move the heavy metal ion to the blood as long as aggregated $A\beta$, that is amyloid, is present. The free $A\beta$ can not penetrate the blood-brain-barrier by passive diffusion, rather it will be degraded by the proteolytic enzymes normally present in the extracellular space. Therefore, free $A\beta$ or a degenerative part thereof may potentially damage the neuronal cells before it is digested to a non-toxic extent. Meanwhile, phanquinone will prevent or inhibit the potential cytotoxic effect of $A\beta$ due to its ability to counteract the reduction of the redox activity $A\beta$ normally produce.

The reduction of the aggregation effected by phanquinone is 50-60% for Cu^{2+} and 10% for Zn^{2+} . It is further shown that clioquinol has the opposite tendency. Clioquinol reduces the Zn^{2+} -induced aggregation of $A\beta(1-40)$ by more than 60%, whereas the Cu^{2+} -induced aggregation was reduced by approximately 30%. In other words, phanquinone is best at reducing the Cu^{2+} induced aggregation of $A\beta(1-40)$ and clioquinol is best at reducing the Zn^{2+} induced aggregation.

As Cu^{2+} and Zn^{2+} naturally are present in the body of a patient having or expected of having Alzheimer's disease, it may be desired to reduce or resolubilize

the aggregation induced by both ions. Thus, administering of clioquinol prior to, together with or subsequent to the administering of phanquinone may be preferred.

The administering of clioquinol is, however, 5 problematic as clioquinol as a side effect causes SMON. Prior studies (see WO 98/06403) indicate a possible influence of clioquinol on vitamin B₁₂.

It is known that clioquinol is excreted through the kidneys as glucuronide or sulphate derivatives 10 (Kotaki H., et al.: Enterohepatic circulation of clioquinol in the rat, *J. Pharmacobiodyn.* 1984 June; 7(6): 420-5 and Jurima M. et al.: Metabolism of 14C-iodochlorhydroxyquinoline in the dog and the rat, *J. Pharmacobiodyn.* 1984 March; 7(3): 164-70), e.g. as the 15 compound methyl(5-chloro-7-iodo-quinolyl-2',3',4'-tri-O-acetyl-glucopyranosid)uronate. For short, this compound is referred to as clioquinol glucuronide in the following.

The detoxification of hydrophobic substances, such 20 as clioquinol, in the body predominantly occurs in the liver. Therefore, it is believed that the clearance of clioquinol happens as follows: Clioquinol is converted to clioquinol glucuronide in the liver. Following the formation, the water soluble clioquinol glucuronide is 25 secreted to the bile. The bile enters the intestine, wherein a major amount of the clioquinol glucuronide is evacuated in the stool. A certain amount of the clioquinol glucuronide is resorbed from the intestine to the blood. The clioquinol glucuronide is filtered from 30 the blood in the kidneys and appears in the terminal urine.

By treating mice with clioquinol and subsequently administering a radioisotope of vitamin B₁₂ ([⁵⁷Co]-cyanocobalamine) it is shown in international patent 35 application, publication No. WO 98/06403 that the

concentration of vitamin B₁₂ in the brain and the liver of the clioquinol-treated mice remains at a normal level, whereas the concentration of the radioisotope of vitamin B₁₂ is decreased in the kidney of such mice compared to the normal level. This finding suggests a metabolism of vitamin B₁₂ being dependent on clioquinol. Further, the finding suggests that the kidneys are the target organs, wherein the clioquinol dependent metabolism occurs.

In order to investigate a possible interaction between clioquinol glucuronide and vitamin B₁₂, an experiment was designed, wherein clioquinol glucuronide and vitamin B₁₂ were mixed in water. The mixture was analyzed by ¹H NMR. The ¹H NMR spectra, see Figs. 5-7, show that some of the resonances of vitamin B₁₂ (corresponding to the benzimidazole moiety) have shifted, and the same is observed for two resonances of the clioquinol glucuronide (corresponding to the quinoline moiety). It is believed by applicant that similar results would be expected using free clioquinol, however clioquinol cannot be dissolved in aqueous solutions for NMR testing.

The results indicate a hydrophobic interaction between clioquinol glucuronide and vitamin B₁₂, possibly between the benzimidazole moiety of the vitamin B₁₂ and the quinoline moiety of clioquinol glucuronide.

Vitamin B₁₂ is normally resorbed actively from the renal plasma after it has been filtered. In that way the body recovers most of the vitamin B₁₂ that would otherwise have been lost in the urine. It has recently been demonstrated that the resorption of vitamin B₁₂ is mediated by the action of the membrane protein megalin (Moestrup S.K. et al. Proc. Natl. Acad. Sci. 1996; 93(16): 8612-7). The megalin is shown to have a strong

affinity towards the binding of a complex formed by vitamin B₁₂ and the transport protein transcobalamin.

Based on new finding reported herein, viz. that vitamin B₁₂ binds to clioquinol glucuronide, it is believed that vitamin B₁₂ does not bind to the megalin protein and/or the transcobalamin when it is already bound to the clioquinol glucuronide. Thus, the resorption of vitamin B₁₂ will fail and the body will suffer from vitamin B₁₂ deficiency after a certain time period of clioquinol administration if the body is not supplied with enough new vitamin B₁₂ through the normal diet.

The only source of vitamin B₁₂ input in man is food, since man cannot synthesize it. Thus, diets low in meat and/or microorganisms will evidently cause vitamin B₁₂ deficiency. If persons are supplied with diets with a too low content of vitamin B₁₂ the administration of clioquinol will worsen the condition as the resorption of B₁₂ is prevented by the competitive binding of clioquinol with megalin. The fact that SMON only was observed in Japan and that the Japanese diet predominantly consisted of vegetables and cereals, especially rice, (Kromhout D, et al : "Food consumption pattern in the 1960s in seven countries", Am.J. Clin. Nutr. 49: 889-894, 1989) may explain why the SMON disease was confined to the Japan.

To counteract side effects of phanquinone and/or clioquinol administration, it may be desired to co-administer trace metals and/or prosthetic groups. A preferred prosthetic group is vitamin B₁₂ (cyanocobalamine).

By way of example, but not limitation, compositions, the use thereof, and methods of the invention can be indicated for: 1) patients diagnosed with Alzheimer's disease at any clinical stage of the

disease, 2) the prevention of Alzheimer's disease in patients with early or prodromal symptoms or signs, and 3) the delay of the onset or evolution or aggravation of the symptoms and signs of Alzheimer's disease. The 5 methods and compositions of the invention will be, for example, useful for the treatment of Alzheimer's disease, the improvement or alleviation of any symptoms and signs of Alzheimer's disease, the improvement of any pathological or laboratory findings of Alzheimer's 10 disease, the delay of the evolution of Alzheimer's disease, the delay of onset of any Alzheimer's disease, symptoms and signs, the prevention of occurrence of Alzheimer's disease, and the prevention of the onset of any of the symptoms and signs of Alzheimer's disease or 15 Parkinson's disease.

The subject, or patient, is an animal, e.g. a mammal, and is preferably human, and can be a fetus, child, or adult.

The dose of phanquinone optimal in vivo for the 20 resolubilisation of A β can be determined by a physician upon conducting routine experiments. An example of such an experiment is the monitoring of soluble A β in the cerebrospinal fluid (CSF) (WO 93/10459, dated May 27, 1993 of University of Melbourne). Beginning with 25 relatively low doses (10-25 mg/day), a physician can monitor the amount of solubilized A β in the patient's CSF. If there is no increase in solubilized A β in response to the phanquinone administration, indicative of resolubilization of zinc-A β aggregates, the dosage 30 can be raised until such an increase is observed. Another example is monitoring the clinical signs and symptoms of the disease by using clinical, behavioural and psychometric observations and measurements.

Prior to administration to humans, the efficacy is 35 preferably shown in animal models. Any animal model for

Alzheimer's disease known in the art can be used.

The pharmaceutical compositions according to the present invention preferably comprise one or more pharmaceutical acceptable carriers and the active
5 constituent(s). The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipients thereof. In a preferred embodiment, the phanquinone and optional further active constituents
10 in the pharmaceutical composition is purified.

It will be appreciated that the amount of phanquinone and optional further active constituents required for said treatment or prevention will vary according to the route of administration, the disorder
15 to be treated, the condition, age, the file history of the subject, and the galenic formulation of the pharmaceutical composition, etc. When treating a patient diagnosed as having Alzheimer's disease, the amount of phanquinone is preferably effective to increase the
20 solubility of A β -aggregations in the cerebrospinal fluid of the patient.

In general, a suitable therapeutically effective amount of phanquinone in the pharmaceutical composition is, for example, 5 to 250 mg, preferably 10 to 50 mg.
25 A suitable amount a compound, or a mixture of compounds, selected from the group comprising antioxidants, μ 1 agonists, anticholinesterase inhibitors, trace metals, prosthetic groups and clioquinol in the pharmaceutical composition is, for example, 5 μ g to 250
30 mg, preferably 0.5 to 1 mg. If clioquinol and vitamin B₁₂ is selected, the amount of clioquinol preferably is effective to the treatment or prevention of Alzheimer's disease and the amount of vitamin B₁₂ is preferably effective to inhibit a detrimental side effect of
35 clioquinol administration. The amounts of clioquinol

and vitamin B₁₂ are preferably 5 mg to 250 mg, most preferred 10 mg to 50 mg and 5 μ to 2 mg, most preferred 0,5 mg to 1 mg, respectively.

5 The actually administered amounts of phanquinone and optional further active constituents such as clioquinol and vitamin B₁₂ may be decided by a supervising physician. If the pharmaceutical composition in addition to phanquinone comprise further active consti-
10 tuents they can be in the same composition for administering in combination concurrently, or in different compositions for administering substantially simultaneously but separately, or sequentially.

Therapeutic formulations include those suitable
15 for parenteral (including intramuscular and intravenous), oral, rectal or intradermal administration, although oral administration is the preferred route. Thus, the pharmaceutical composition may be formulated as tablets, pills, syrups, capsules, suppositories,
20 formulations for transdermal application, powders, especially lyophilized powders for reconstitution with a carrier for intravenous administration, etc.

The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is
25 administered. The carriers in the pharmaceutical composition may comprise a binder, such as microcrystalline cellulose, polyvinylpyrrolidone (polyvidone or povidone), gum tragacanth, gelatine, starch, lactose or lactose monohydrate; a disintegrat-
30 ing agent, such as alginic acid, maize starch and the like; a lubricant or surfactant, such as magnesium stearate, or sodium lauryl sulphate; a glidant, such as colloidal silicon dioxide; a sweetening agent, such as sucrose or saccharin; and/or a flavouring agent, such
35 as peppermint, methyl salicylate, or orange flavouring.

Therapeutic formulations suitable for oral administration, e.g. tablets and pills, may be obtained by compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by mixing the constituent(s), and compressing this mixture in a suitable apparatus into tablets having a suitable size. Prior to the mixing, the phanquinone may be mixed with a binder, a lubricant, an inert diluent and/or a disintegrating agent and the further optionally present constituents may be mixed with a diluent, a lubricant and/or a surfactant.

In a preferred embodiment, free-flowing phanquinone powder is mixed with a binder, such as microcrystalline cellulose, and a surfactant, such as sodium lauryl sulphate, until a homogeneous mixture is obtained. Subsequently, another binder, such as polyvidone, is transferred to the mixture under stirring. When a uniform distribution is obtained an aqueous solution of vitamin B₁₂ is added under constant stirring. This mixture is passed through granulating sieves and dried by desiccation before compression into tablets in a standard compressing apparatus.

In a second preferred embodiment, free-flowing phanquinone powder is mixed with surfactants and/or emulsifying agents, such as Sapamine® (N-(4'-stearoyl amino phenyl)-trimethylammonium methyl sulphuric acid) and lactose monohydrate until a uniform distribution of the constituents is obtained. A second preparation containing a disintegrating agent, such as maize starch, is added to the phanquinone mixture under continuous stirring. Such a second preparation may be prepared by adding excess boiling water to a maize starch suspended in cold water. The final mixture is granulated and dried as above and mixed with maize starch and magnesium stearate and finally compressed

into tablets in a standard apparatus.

A tablet may be coated or uncoated. An uncoated tablet may be scored. A coated tablet may be coated with sugar, shellac, film or other enteric coating agents.

Therapeutical formulations suitable for parenteral administration include sterile solutions or suspensions of the active constituents. An aqueous or oily carrier may be used. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soy bean oil, mineral oil, sesame oil and the like. Formulations for parenteral administration also include a lyophilized powder comprising phanquinone and optionally further active constituents that is to be reconstituted by dissolving in a pharmaceutically acceptable carrier that dissolves the active constituents, e.g. an aqueous solution of carboxymethylcellulose and lauryl sulphate.

When the pharmaceutical composition is a capsule, it may contain a liquid carrier, such as a fatty oil, e.g. cacao butter.

Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides.

In yet another embodiment, the therapeutic compound can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer,

supra; Sefton, CRC Crit. Ref. Biomed. Eng. 14: 201 (1987); Buchwald et al., Surgery 88: 507 (1980); Saudek et al., N. Engl. J. Med. 321: 574 (1989)). In another embodiment, polymeric materials can be used (see 5 Medical Applications of Controlled Release, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J. Macromol. Sci. Rev. 10 Macromol. Chem. 23: 61 (1983); see also Levy et al., Science 228: 190 (1985); During et al., Ann. neurol. 25: 351 (1989); Howard et al., J. Neurosurg. 71: 105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of the 15 therapeutic target, i.e., the central nervous system, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled Release, *supra*, vol. 2, pp. 115-138 (1984)).

Other controlled release systems are discussed in 20 the review by Langer (Science 249: 1527-1533 (1990)).

In one embodiment of the pharmaceutical composition according to the invention, phanquinone and the further active constituents, are comprised as separate pharmaceutical entities. By way of example, one entity 25 may comprise phanquinone and clioquinol and another entity may comprise vitamin B₁₂. The two entities, may be administered simultaneously or sequentially. For example, the entity comprising phanquinone and clioquinol can be administered, followed by vitamin B₁₂ 30 administered within a day, week, or month of clioquinol and phanquinone administration. If the two entities are administered sequentially, the entity comprising phanquinone and clioquinol is preferably administered for one to three weeks followed by a wash out period of 35 one to four weeks during which the entity comprising

vitamin B₁₂ is administered, but not the entity comprising clioquinol. After the wash out period, the treatment can be repeated.

5 The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form
10 described by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

15 Other features and advantages of the invention will be apparent from the following examples, which in conjunction with the accompanying drawings illustrate by way of example the principles of the invention.

20

EXAMPLES

Example 1

Effect of phanquinone on A β (25-35) dose-response in PC12 cells.

25 A β (25-35) was delivered by Bachem (CH) or Sigma (USA) and dissolved in phosphate buffered saline (PBS) at pH 7.4, 2 hours prior to application. The neurotoxicity of A β is located in the sequence between amino acid residues 25 and 35 (A β (25-35)) and a deca-
30 peptide encompassing this region induces neural cell death equally potent as full length A β (1-40) (Yankner, Duffy LK, Kirschner DA: Neurotrophic and neurotoxic effects of amyloid β protein: reversal by tachykinin neuropeptides. *Science* 1990;250(4978):279-282).

35 Rat PC12 pheochromocytoma cells were grown in

Dulbecco's modified Eagle's medium (DMEM) containing 1% penicillin-streptomycin, 5% fetal calf serum and 10% horse serum in humidified incubator with 5% CO₂.

PC12 cells were plated on 96-wells microtiter plates in 100 μ l of the appropriate medium. After 24 hours the indicated concentrations of A β (25-35) peptide was added alone or together with phanquinone in the designated concentrations. Incubation continued for 24 hours. Following incubation, MTT reduction was measured using a commercially available assay according to the manufacturer's (Boehringer Mannheim) instructions. Assay values obtained by vehicle alone were defined as 100%.

MTT is a substrate for intracellular and plasma membrane oxidoreductases and has been widely used to measure reductions of cell redox activity. Reduction of the cell redox activity has been found to be an early indicator of A β mediated cell death (Shearman MS, Ragan CI, Iversen LL: Inhibition of PC12 cell redox activity is a specific, early indicator of the mechanism of β -amyloid-mediated cell death, *Proc. Natl. Acad. Sci. USA* 1994;91(4):1470-1474).

To test the effect of phanquinone on A β (25-35) induced toxicity in PC12 cells, PC12 cells were exposed to A β (25-35) peptide concentrations ranging from 0 to 10 μ M. In the absence of phanquinone (i.e. only vehicle added) A β (25-35) produced a dose-dependent inhibition of MTT reduction (Fig. 1, filled-in circles). Concentrations of A β (25-35) as low as 0,01 μ M produced a significant reduction and at a concentration of A β (25-35) on or above 0,1 μ M the MTT reduction was reduced to a maximum level of about 50%.

In the presence of 10 μ g phanquinone per ml the toxic effect of A β (25-35) was virtually abolished (Fig. 1, open circles). Even at concentrations of A β as high

as 1 μ M, the presence of phanquinone completely counteracted the toxic effect of A β . Only at the highest concentration of A β (10 μ M) there was a slight inhibition of MTT reduction by A β . This inhibition was, 5 however moderate, only around 10% as compared to approximately 50% in the absence of phanquinone.

Example 2

Effect of phanquinone in A β (25-35) induced toxic-
10 ity in PC12 cells.

In this study the same materials and methods as in example 1 were used, except that the PC12 cells were exposed to a fixed A β concentration of 1 μ M, whereas the concentration of phanquinone was varied between 0
15 and 10 μ g/ml. In the absence on phanquinone the A β (25-35) resulted in approximately 40% inhibition of MTT reduction (Fig. 2, open circles). This inhibition was maintained in the presence of up to 0,1 μ g phanquinone per ml. Increasing the concentration of phanquinone
20 above 0,1 μ g/ml resulted in drastic reduced toxic effect of A β (25-35). At concentrations of phanquinone of 1 μ g/ml and above, the toxic effect of A β (25-35) was completely abolished.

25 Example 3

Effect of phanquinone and clioquinol on Zn²⁺- and Cu²⁺-induced A β aggregation.

A 5 mg/ml stock solution of A β (1-40) (delivered from Bachem (CH)) was freshly prepared before each
30 experiment by dissolving the lyophilised peptide in 0,01 M HCl, followed by subsequent dilution 1:1 with 0,01 M NaOH to yield a neutral pH. Aliquot of A β (1-40) were diluted in PBS (pH 7,4) to 100 μ M and incubated at a total volume of 30 μ l for 24 hours at room tempera-
35 ture. For co-incubation experiments, the indicated

concentrations of metal ions and/or aliquot of test compounds were added. The test compounds were added to a final molar concentration of 10 μ g/ml.

The amyloid formation was quantified by a thioflavin T fluorometric assay. Thioflavin binds specifically to amyloid and this introduces a shift in its emission spectrum and a fluorescent signal proportional to the amount of amyloid is formed. After incubation, A β (1-40) peptides were added to PBS (pH 6,0) and 3 μ M thioflavin T in a final volume of 1 ml. Fluorescence was monitored at excitation 454 nm and emission 482 nm using a Fluoroscan II fluorometer (Molecular devices, UK). A time scan of fluorescence was performed and three values after the decay reached a plateau (around 5 minutes) were averaged after subtracting the background fluorescence of 3 μ M thioflavin T. For co-incubation experiments, fluorescence of test compound alone was determined. Samples were run in triplicate. The mean \pm SD for the typical experiment is shown in the figures (Fig. 3 and 4).

Phanquinone and clioquinol were tested for their ability to prevent the aggregation of A β (1-40) into amyloid structures. Clioquinol is a known compound in the treatment of Alzheimer's disease (see international publication number WO 98/06403) and was originally tested in the present experiment for comparison purposes.

It was studied whether the two compounds had any effect on metal-ion catalysed A β aggregation, especially the aggregation caused by Zn²⁺ and Cu²⁺. The results of the experiments are shown on Fig. 3 and 4.

In the Figures 3 and 4 it is revealed that Cu²⁺ and Zn²⁺ to a lesser extent, increase the aggregation of A β into amyloid structures relative to the spontaneous aggregation. In the presence of phanquinone and

cliöquinol at concentrations of 10 µg/ml, the metal ion induced aggregation of Aβ was significantly reduced.

At the tested concentration of 10 µg/ml phanquinone reduced the Cu²⁺-induced aggregation by 50-60%,
5 while the Zn²⁺-induced aggregation was only modest inhibited by approximately 10%. Unexpectedly, clioquinol showed the opposite tendency. Clioquinol reduced

the Zn²⁺-induced aggregation of Aβ(1-40) by more than
10 60%, whereas the Cu²⁺-catalysed aggregation was reduced by approximately 30%. A pharmaceutical composition comprising phanquinone in combination with clioquinol may thus have a more widely usage than a pharmaceutical composition comprising one of the compounds alone.

15


Example 4

In this example, a metabolite of clioquinol was synthesized.

It is known that the clioquinol is excreted
20 through the kidneys as glucuronide derivatives of clioquinol (Kotaki H., et al.: "Enterohepatic circulation of clioquinol in the rat", J. Pharmacobiodyn. 1984 June; 7(6): 420-5 and Jurima M. et al.: "Metabolism of 14C-iodochlorhydroxyquinoline in the dog and the rat",
25 J. Pharmacobiodyn. 1984 March; 7(3): 164-70). The transformation of clioquinol to the corresponding glucuronide presumable takes place in the liver. Following the formation of clioquinol glucuronide in the liver it is eventually transferred to the kidneys for excretion
30 in the urine.

A glucuronide derivative of clioquinol in the form of methyl(5-chloro-7-iodo-quinolyl-2',3',4'-tri-O-acetyl-glycopyranosid)uronate was prepared according to the following reaction scheme:

23



A mixture of 5-chloro-8-hydroxy-7-iodo-quinoline (50 mg, 0.164 mmol), methyl 1-bromo-1-deoxy-2,3,4-tri-O-acetyl-D-glucopyranosiduronate (65 mg, 0.164 mmol), $\text{CaSO}_4 \cdot \text{H}_2\text{O}$ (35 mg) and pyridine (1.5 ml) was stirred at room temperature for 20 min. Freshly prepared Ag_2CO_3 (35 mg) was added to the reaction mixture and the suspended solution was stirred at room temperature for 20 hours in the dark. Subsequently, the reaction product was deacetylated by 1 N aqueous NaOH.

10 The reaction mixture was diluted with CH_2Cl_2 (10 ml), filtered and the solvent evaporated under reduced pressure. The above product was purified by flash chromatography (TLC: $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 99/1, eluent: $\text{CH}_2\text{Cl}_2/-\text{MeOH}$ 99.5/0.5).

15 NMR (400MHz, CDCl_3) 2.04 (s, 3H, Ac), 2.09 (s, 3H, Ac), 2.13 (s, 3H, Ac), 3.68 (s, 3H, Me), 3.99 (d, 1H, 5'-H), 5.40-5.52 (m, 3H, 2', -3', -4'-H), 6.29 (d, 1H, 1'-H), 7.56 (m, 1H, 3H), 7.99 (s, 1H, 6-H), 8.52 (d, 1H, 4-H), 8.93 (s, 1H, 2-H).

20 This compound is referred to as clioquinol glucuronide in the following.

Example 5

The interaction of vitamin B_{12} with clioquinol glucuronide as prepared in example 4, was studied using nuclear magnetic resonance (NMR) spectroscopy.

As the clioquinol glucuronide is soluble in water, the study was undertaken in buffered water at pH=6.5. Three different solutions were prepared and their ^1H NMR spectra were recorded in a DRX 400 MHz spectrophotometer at 20°C. Solution a) contained free vitamin B_{12} (hydroxycobalamin) in a concentration of 2.6 mM. Solution b) contained a mixture of 2.6 mM vitamin B_{12} and 10 mM clioquinol glucuronide (mole ratio of about 1:4). Solution c) contained 10 mM clioquinol glucu-

ronide.

In Fig. 5 the spectra of the three solutions are presented for the aromatic region (5.5-9.8 ppm). The differences are quite small but obvious in the expansion shown of Fig. 6 and Fig. 7, respectively. Some of the resonances of vitamin B₁₂ (corresponding to the benzimidazole moiety) have shifted (see Fig. 6), and the same is observed for two resonances of the clioquinol glucuronide (corresponding to the quinoline moiety) (see Fig. 7).

The results suggest an interaction between clioquinol glucuronide and vitamin B₁₂, possibly of a hydrophobic nature between the benzimidazole moiety of the vitamin B₁₂ and the quinoline moiety of clioquinol glucuronide.

The hydrophobic binding of vitamin B₁₂ to clioquinol glucuronide is believed to cause the vitamin B₁₂ to be excreted from the body together with clioquinol glucuronide, thus preventing resorption of vitamin B₁₂, which would eventually lead to a vitamin B₁₂ deficiency. Therefore, vitamin B₁₂ deficiency is believed to be, at least to some extent, the underlying cause of SMON. In consequence, whenever clioquinol is administered it should be ensured that the level of vitamin B₁₂ in the treated subject is sufficient for avoiding deficiency. This may be accomplished by co-administering of clioquinol and vitamin B₁₂.

Example 6

Preparation of a pharmaceutical composition comprising phanquinone.

250 g of phanquinone was mixed with 200 g sapsamine[®] (N-(4'-stearoyl amino-phenyl)-trimethylammonium methyl sulphuric acid) and 1025 g lactose mono-hydrate for a period of 5 minutes. 300 g of boiling water was

added in one go to a mixture of 100 g maize starch in 100 g cold water. The maize suspension, cooled to 40°C was added to the phanquinone containing powder mixture under continuous stirring. The mixture was granulated using a 2.5 mm sieve and desiccated for 18 hours at 40°C. The dry granules were mixed with 400 g maize starch and 20 g magnesium stearate. The final mixture was formulated into tablets having a diameter of 8.0 mm and a weight of 200 mg.

10

Example 7

Preparation of a pharmaceutical composition comprising phanquinone, clioquinol and vitamin B₁₂.

250 g phanquinone and 250 g of clioquinol was mixed with 200 g sapamine[®] (N-(4'-stearoyl amino-phenyl)-trimethylammonium methyl sulphuric acid) and 1025 g lactose mono-hydrate for a period of 5 minutes. 300 g of boiling water was added in one go to a mixture of 100 g maize starch in 100 g cold water. The maize suspension, cooled to 40°C was added to the phanquinone and clioquinol containing powder mixture under continuous stirring. Subsequently, an aqueous solution of 5 g vitamin B₁₂ was added. The mixture was granulated using a 2.5 mm sieve and desiccated for 18 hours at 40°C. The dry granules were mixed with 400 g maize starch and 20 g magnesium stearate. The final mixture was formulated into tablets having a diameter of 8.0 mm and a weight of 200 mg.

Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties.

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention, and all such

modifications, as would be obvious to a person skilled in the art, are intended to be included in the scope of the following claims.